



## SUPPLEMENTAL LABEL

### POAST<sup>®</sup> HERBICIDE

EPA Reg. No. 7969-58-51036

AD041101

All applicable directions, restrictions and precautions on the EPA-registered label are to be followed. This labeling must be in the possession of the user at the time of application.

## For Tall Fescue Growth Suppression to Aid in the Establishment of Forbs or Other Grasses

### Directions For Use

It is a violation of federal law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your state or tribe, consult the agency responsible for pesticide regulation.

### Application Instructions

Apply **Poast** to actively growing tall fescue after it has 4-6 inches of new growth, before the emergence of seedheads and before conifer bud break. Applications made from July 1 to mid-August may be less effective, especially if day temperatures reach 90° F. Tall fescue must be 1-year old before the first application of **Poast**.

Adequate coverage of the leaf surface is necessary for absorption of the herbicide. Thus, for optimum control, do not mow tall fescue turf for 30 days before or 14 days after applying **Poast**.

**Rate:** Apply 1.0-1.25 pints of **Poast** per acre. For greater fescue suppression, up to 2.5 pints of **Poast** per acre can be used. Because of environmental differences at application and growth differences of tall fescue, control may exceed or fall short of that desired. Begin treating crops with **Poast** at the minimum recommended rate and adjust rates as local conditions and experience dictate. Additional applications may be made if extended growth suppression is desired.

### Additives

Always add two pints of oil concentrate per acre.

### Restrictions and Limitations

**Poast herbicide** should only be applied as a directed spray to the grove floor for bearing pistachio trees.

Avoid all direct or indirect contact with any desired grass crop unless otherwise recommended on the **Poast** label.

Do not apply to grasses or crops under stress such as stress due to lack of moisture, hail damage, flooding, herbicide injury, mechanical injury, or widely fluctuating temperatures, as unsatisfactory control may result. Irrigation may be needed.

Do not apply to crops that show injury (leaf phytotoxicity or plant stunting) produced by any other prior herbicide applications, because this injury may be enhanced or prolonged.

Physical incompatibility, reduced weed control, or crop injury may result from mixing **Poast** with other pesticides (fungicides, herbicides, insecticides, or miticides), additives or fertilizers. MICRO FLO COMPANY, LLC does not recommend using tank mixes other than those listed on MICRO FLO COMPANY, LLC labeling. Local agricultural authorities may be a source of information when using other than MICRO FLO COMPANY, LLC recommended tank mixes.

*Poast is a registered trademark of BASF.*

Manufactured By:  
**Micro Flo Company LLC**  
P.O. Box 772099  
Memphis, Tn. 38117